

DOUBLE COMPACT OBJECTS AS LOW-FREQUENCY GRAVITATIONAL WAVE SOURCES

KRZYSZTOF BELCZYNSKI^{1,2}, MATTHEW BENACQUISTA³, AND TOMASZ BULIK^{2,4}

¹ Los Alamos National Laboratory, CCS-2 Group (Oppenheimer Fellow)

² Astronomical Observatory, University of Warsaw, Al. Ujazdowskie 4, 00-478 Warsaw, Poland

³ Center for Gravitational Wave Astronomy, University of Texas at Brownsville, Brownsville, TX 78520, USA

⁴ Nicolaus Copernicus Astronomical Center, Warsaw, Poland

kbelczyn@nmsu.edu, benacquista@phys.utb.edu, tb@astrouw.edu.pl

Draft version February 26, 2010

ABSTRACT

We study the galactic population of double compact objects (NS-NS, BH-NS, BH-BH binaries) to investigate the number (if any) of these systems that can potentially be detected with LISA at low gravitational-wave frequencies. We calculate the Galactic numbers and physical properties of these binaries and show their relative contribution from the disk, bulge and halo. It is found that, although the formation efficiency of double compact objects is very small, a few thousand of such systems should be detectable with LISA depending on the instrument low-frequency sensitivity and the efficiency of data analysis techniques for identifying highly eccentric systems. The vast majority of these binaries will add up to foreground noise arising from the galactic double white dwarf binaries, and only a very few ($\sim 1 - 10$) are predicted to stand out above the noise and be resolvable. Detailed population synthesis, combined with an updated Galactic model and a realistic LISA signal simulator are used to obtain the results. We outline the prospects of the science that could be done with future LISA observations of double compact objects. In particular, we note that LISA observations could be used to determine the fate of massive binaries that are the most likely progenitors of BH-BH systems and shed some light on the common envelope evolution involved in the formation of double compact objects. Finally, the galactic double compact object populations presented here may readily serve in LISA data mock challenges.^a

^aThe data can be obtained upon request from Chris Belczynski.

Subject headings: binaries: close — stars: evolution, neutron — gravitation

1. INTRODUCTION

The *Laser Interferometer Space Antenna* (LISA) is a next generation instrument to search for and observe gravitational radiation (e.g., Hughes 2006). A system of three satellites (5 million kilometers apart), orbiting the Sun, will form an interferometer sensitive to low frequency gravitational radiation (GR) ($\sim 5 \times 10^{-5} - 1$ Hz). The main sources of GR at these low frequencies are inspirals of super-massive black holes in the center of merging galaxies, extreme mass ratio inspirals of stars and stellar mass black holes into the super-massive black holes, and also nearby (mostly Galactic) compact binaries. Double white dwarf binaries (WD-WD) are the largest population of Galactic systems that are expected to produce a confusion-limited noise in the detector, with several thousand of the louder systems being potentially resolved. A number of studies have concentrated on studies of double white dwarfs in the context of low-frequency LISA observations (e.g., Hils, Bender & Webbink 1990; Farmer & Phinney 2003; Nelemans, Yungelson & Portegies Zwart 2004; Ruiter et al. 2009a,b). Here, we focus on the other, and much less studied group of compact systems; double neutron star binaries (NS-NS), black hole neutron stars (BH-NS), and double black holes (BH-BH). Although much less common than double white dwarfs, these systems may produce much stronger GR signals, since these compact objects are much denser and more massive than white dwarfs.

Out of all double compact objects only NS-NS have so far been observed. A handful of NS-NS were discov-

ered in radio surveys (e.g., Lorimer 2005). Seven systems are found in the Galactic field, while one belongs to the Galactic globular cluster M15 (PSR B2127-11C). Out of the seven, three are wide systems with merger time over Hubble time (PSR J1518+4804, PSR J1811-1736, PSR J1829+2456). The remaining four constitute the close NS-NS population with relatively short ($\sim 0.1 - 3$ Gyr) merger times (PSR B1913+16, PSR B1534+12, PSR J1756-2251, PSR J0737-3039). Also one additional field system (PSR J1906+0746) is considered as a very probable close NS-NS (merger time ~ 0.3 Gyr) binary (Kasian 2008), although the possibility that this is a NS-WD binary can not yet be excluded.

Mergers of all types of double compact objects are expected to be the prime candidates for high-frequency ground based gravitational-wave interferometers like LIGO or VIRGO, while mergers of NS-NS, BH-NS and BH-NS were proposed as potential progenitors of short-hard Gamma-ray bursts (Paczynski 1986). Double compact object binaries were studied extensively over last 2 decades, bringing new developments and understanding of their formation in particular in context of population synthesis studies (e.g., Tutukov & Yungelson 1994; Lipunov, Postnov & Prokhorov 1997; Portegies Zwart & Yungelson 1998; Bethe & Brown 1998; Belczynski & Bulik 1999; Fryer, Woosley & Hartmann 1999; Bloom, Sigurdsson & Pols 1999; Nelemans, Yungelson & Portegies Zwart 2001; Hurley, Tout & Pols 2002; Voss & Tauris 2003; Dewi & Pols 2003; Pfahl, Podsiadlowski & Rappaport 2005).

Merger rates were recently presented and discussed by Kim, Kalogera & Lorimer (2006; empirical estimates) and Belczynski et al. (2007; population synthesis).

In this study, we analyze the entire Galactic population (disk, bulge and halo) of double compact objects. Using population synthesis methods we predict numbers and physical properties of double compact objects, then we calculate their spatial distribution and finally estimate the low frequency gravitational radiation signal that may arise from these binaries. In §2 we describe our method of calculation, in §3 we present results, in §4 we compare our findings with previous work, and finally in §5 we present our conclusions.

2. MODELING

2.1. Population Synthesis

We have used the population synthesis code **StarTrack** to calculate the numbers and properties of double compact objects. The full description of the code can be found in Belczynski, Kalogera & Bulik (2002) and Belczynski et al. (2008). The code utilizes a set of stellar models (Hurley et al. 2000) that allow for evolution of stars at different metallicities. The compact object formation follows self-consistently from the stellar models, extended to the formation of the FeNi core (Timmes, Woosley & Weaver 1996). During the core collapse the fall back and direct black hole formation is accounted for (Fryer & Kalogera 2001) and the newly born compact objects receive natal kicks (Hobbs et al. 2005). Formation of low mass neutron stars through electron capture supernovae is also accounted for (e.g., Podsiadlowski et al. 2004). Binary interactions are treated in detail and the various processes were calibrated using either results of detailed evolutionary calculations (e.g., Wellstein, Langer & Braun 2001 for mass transfer sequences) or specific sets of observations (e.g., Levine, Rappaport & Zojcheski 2000 for tidal interactions).

All of our population synthesis calculations implement the standard evolutionary model presented in Belczynski et al. (2008). We employ our standard model to evolve three different Galactic populations: disk, bulge and halo. The dynamical evolution of stars in these components is described in the following section. Here we describe the evolutionary aspect of the stars in each component. The disk is assumed to have a stellar mass of $4.0 \times 10^{10} M_{\odot}$, stars have solar metallicity $Z = 0.02$ and we assume the age of the population to be 10 Gyr with constant star formation throughout the disk lifetime (i.e., star formation rate is $4 M_{\odot} \text{ yr}^{-1}$). The bulge is assumed to have a stellar mass of $1.1 \times 10^{10} M_{\odot}$, stars also have solar metallicity $Z = 0.02$ and we assume the age of the population to be 10 Gyr with a burst of star formation that lasted throughout first Gyr of bulge evolution (i.e., all the stars were born between 9 and 10 Gyr ago; a uniform distribution of birth times was adopted for all the stars in this time interval). The halo is assumed to have a stellar mass of $0.1 \times 10^{10} M_{\odot}$, the stars have sub-solar metallicity $Z = 0.001$ and we assume the age of the population to be 13 Gyr with a burst star formation at the very beginning of halo evolution (i.e., all the stars were born 13 Gyr ago in a delta function-like burst).

We assume that in each population the binary fraction is

50% (2/3 of stars are in binaries, and 1/3 are single stars). We do not consider any dynamical interactions between stars; i.e., evolution of stars in globular clusters is not accounted for and we evolve only field populations. However, we note that despite their relative small stellar mass content globular clusters may contribute significantly to formation of BH-BH binaries (Kulkarni, Hut & McMillan 1993; Portegies Zwart & McMillan 2000; Gultekin, Miller & Hamilton 2004; O’Leary et al. 2006; Sadowski et al. 2008). On the other hand, formation of double compact object systems with neutron stars was found to be rather inefficient in globular cluster environments (e.g., Phinney 1991; Grindlay, Portegies Zwart, & McMillan 2006; Ivanova et al. 2008). The initial mass of single stars and binary primaries (more massive component) are drawn from the three component broken power-law Initial Mass Function (IMF). We employ the IMF from Kroupa, Tout & Gilmore (1993) and Kroupa & Weidner (2003) with slope of -1.3 ($0.08 < M < 0.5 M_{\odot}$), -2.2 ($0.5 < M < 1.0 M_{\odot}$), -2.7 ($1.0 < M < 150 M_{\odot}$). The binary secondaries are drawn through flat mass ratio distribution within the entire mass range ($0.08 < M < 150 M_{\odot}$). We follow only the evolution of massive binaries as only these can form double compact object binaries. We require that the mass of the primary and secondary are the following: $M_1 > 6 M_{\odot}$ and $M_2 > 4 M_{\odot}$. Even though it is expected that only stars more massive than $\sim 8 M_{\odot}$ can form compact objects (i.e., neutron stars and black holes), the binary evolution (through mass transfer and rejuvenation) can alter such a simple picture. The binary initial separations are drawn from distribution that is flat in logarithm (Abt 1983) with maximum separations reaching $10^5 R_{\odot}$. Binaries are allowed to be eccentric, and we use thermal distribution for initial eccentricities (i.e., $2e$; Heggie 1975; Duquennoy & Mayor 1991).

It was noted by Belczynski et al. (2007) that many progenitors of double compact object binaries evolve through a common envelope phase while a donor star is crossing Hertzsprung gap. Such a star does not have a well developed core-envelope structure (e.g., Taam & Sandquist 2000) and once inspiral in a common envelope is started it may very likely lead to a merger whether there is enough orbital energy to expel the envelope or not. If such mergers are taken into account they lead to a significant decrease in the formation of close double compact objects. To estimate the impact of this uncertainty on the LISA signal we perform two calculations. In one (Model A) we allow for survival in the case of a common envelope with a Hertzsprung gap donor (i.e., standard energy balance is tested to check for system survival; Webbink 1984), while in another (model B) we assume that all such common envelope events lead to a merger aborting further binary evolution and thus depleting the population of double compact objects.

2.2. Galactic Model

The binaries are distributed at birth throughout the Galaxy according to stellar density models. We assume that the densities are independent of time. For the bulge, we choose a spherical density with a normal distribution in the radial coordinate and a cut-off radius of 3.5 kpc, so

$$\rho_b \propto e^{-(r/r_{\text{ob}})^2}, \quad (1)$$

where $r_{0b} = 500$ pc (Nelemans et al. 2004). The disk population is assumed to be axially symmetric with cylindrical radius and vertical height distribution given by a double exponential:

$$\rho_d \propto e^{-R/R_{0d}} e^{-|z|/z_{0d}}, \quad (2)$$

with $R_{0d} = 2.5$ kpc and $z_{0d} = 200$ pc. Finally, we distribute the halo binaries according to a simplistic spherical model with:

$$\rho_h \propto \left(1 + \frac{r}{a_0}\right)^{-3.5}, \quad (3)$$

with $a_0 = 3.5$ kpc. The disk and bulge systems are given initial rotational velocities in the plane of the disk, and the halo systems are assumed to have circular orbits corresponding to their initial positions. All systems propagate through a Galactic potential and their trajectories change due to the kicks received at the birth of each compact object.

The Galactic gravitational potential is the sum of the bulge, disk and halo potentials. The disk and bulge are described the Miyamoto and Nagai (1975) potentials:

$$\Phi(R, z) = \frac{GM_i}{\sqrt{R^2 + (a_i + \sqrt{z^2 + b_i})}} \quad (4)$$

where $R = \sqrt{x^2 + y^2}$, the index i corresponds to bulge and disk, a_i and b_i are the parameters, and M_i is the mass. The halo potential is assumed to be

$$\Phi(r) = -\frac{GM_h}{r_c} \left[\frac{1}{2} \ln \left(1 + \frac{r^2}{r_c^2} \right) + \frac{r_c}{r} \operatorname{atan} \left(\frac{r}{r_c} \right) \right] \quad (5)$$

where r_c is the core radius, and M_h is the parameter describing the mass of the halo (Paczynski 1990). The mass of such halo is divergent so we introduce a cutoff radius r_{cut} beyond which the halo density falls to zero, and the potential is $\Phi(r) \propto r^{-1}$. The bulge is described by $M_1 = 1.12 \times 10^{10} M_\odot$, $a_1 = 0$ kpc, and $b_1 = 0.277$ kpc, and the disk by $M_2 = 8.78 \times 10^{10} M_\odot$, $a_2 = 4.2$ kpc, and $b_2 = 0.198$ kpc. For the halo potential we use $r_c = 6.0$ kpc, $r_{cut} = 100$ kpc, and $M_h = 5.0 \times 10^{10} M_\odot$. We note that the masses of the potentials differ from the masses of the stellar components in Section 2.1 because we include non-stellar matter and dark matter in the potentials.

2.3. LISA Signal Simulator

We calculate the expected signal in LISA as the Michelson signal from an equal-arm interferometer using the long wavelength approximation for frequencies below 3 mHz (Cutler 1998; Benacquista, DeGoes & Lunder 2004) and the rigid adiabatic approximation (Rubbo, Cornish & Poujade 2004) above this frequency. The gravitational waveform for an eccentric binary with angular frequency $\omega = 2\pi/P_{orb}$ is calculated in the quadrupole approximation (Peters & Mathews 1963; Peters 1964). We use the form given in Pierro et al. (2001):

$$h_+ = \frac{1}{\sqrt{2}} \left[(1 + \cos^2 \vartheta) [2h_{xy} \sin 2\varphi + h_{x-y} \cos 2\varphi] - (1 - \cos^2 \vartheta) h_{x+y} \right] \quad (6)$$

$$h_\times = \frac{\cos \vartheta}{\sqrt{2}} [2h_{xy} \cos 2\varphi - h_{x-y} \sin 2\varphi] \quad (7)$$

where ϑ and φ are the usual spherical angular coordinates in the frame centered on the binary with the x -axis along the periastron. The metric components appear as:

$$h_{xy} = \sum_{n=1}^{\infty} h_{xy}^{(n)} \sin(n\omega t + n\phi_0) \quad (8)$$

$$h_{x\pm y} = \sum_{n=1}^{\infty} h_{x\pm y}^{(n)} \cos(n\omega t + n\phi_0) \quad (9)$$

where ϕ_0 is the phase of the binary at $t = 0$ and

$$h_{xy}^{(n)} = h_0 n \sqrt{1 - e^2} [J_{n-2}(ne) + J_{n+2}(ne) - 2J_n(ne)] \quad (10)$$

$$h_{x-y}^{(n)} = 2h_0 \{n [J_{n-2}(ne) - J_{n+2}(ne)] - 2ne [J_{n-1}(ne) - J_{n+1}(ne)] + 2J_n(ne)\} \quad (11)$$

$$h_{x+y}^{(n)} = -4h_0 J_n(ne) \quad (12)$$

where $J_n(x)$ are Bessel functions. The overall amplitude of the gravitational wave is proportional to:

$$h_0 = \frac{2G^{5/3} \omega^{2/3} M_{chirp}^{5/3}}{rc^4}, \quad (13)$$

where r is the luminosity distance from the earth to the binary. We assign each binary an arbitrary orientation, and so the contribution of each binary to the LISA signal is described by 11 parameters. The masses (M_1 and M_2), luminosity distance (r), and orbital frequency (f_{orb}) are used to construct the overall amplitude and the relative amplitudes of the harmonics are obtained from the eccentricity (e). The sky location is obtained from the final positions (θ_s and ϕ_s) of the binaries after propagation through the Galactic potential. The arbitrary orientation includes the initial phase (ϕ_0), the argument of the periastron (γ), and the direction angles of the angular momentum vector (θ_l and ϕ_l). We compute the one-year time domain signal for each harmonic in the waveform up to a frequency of 30 mHz. We have not included periastron precession, although this may prove to be important for longer observation times.

For $e = 0$, all the power in the waveform is concentrated at the $n = 2$ harmonic. At non-zero eccentricity, the power in this waveform is spread out over several harmonics of the orbital frequency. If we define $n_{max}(e)$ to be the harmonic with maximum power for eccentricity e , then for $e \gtrsim 0.8$, $n_{max}(e)$ can be approximated by

$$n_{max} \simeq \frac{1.6}{(1 - e)^{1.5}}. \quad (14)$$

The factor of 1.6 arises from maximizing $g(n, e)$ from Peters & Mathews (1963), but can be understood by realizing that the speeds of the components are larger at pericenter than for the corresponding circular orbit. This can be understood by taking into account that the gravitational waves must emit close to the frequency of a circular orbit with radius equal to the periastron, r_{peri} , which is given by:

$$f_{max} \propto r_{peri}^{-3/2} \propto (1 - e)^{-1.5} f_{orb} \propto n_{max} f_{orb}. \quad (15)$$

Therefore, binaries with f_{orb} below the low-frequency limit (f_{crit}) for LISA sensitivity may still be observable by LISA if $n_{\text{max}} f_{\text{orb}} > f_{\text{crit}}$. We use the following approximate criterion to separate potential LISA sources from the entire double compact object population

$$P_{\text{orb}} < 1.6(1 - e)^{-1.5} f_{\text{crit}}^{-1} \quad (16)$$

where $f_{\text{crit}} = 5 \times 10^{-5}$ Hz.

The LISA noise is simulated by assuming the power spectral density of the noise is made up of position (or shot) noise given by (Cornish, 2001):

$$S_{\text{np}} = 4.8 \times 10^{-42} \text{ Hz}^{-1}, \quad (17)$$

and an acceleration noise (converted to strain) given by:

$$S_{\text{na}} = 2.3 \times 10^{-40} \left(\frac{10^{-3} \text{ Hz}}{f} \right)^4 \text{ Hz}^{-1}. \quad (18)$$

These separate components are combined according to:

$$S_{\text{n}} = 4S_{\text{np}} + 8S_{\text{na}} (1 + \cos^2(f/f_*)), \quad (19)$$

where $f_* = c/2\pi L$ with the armlength of LISA taken to be $L = 5 \times 10^9$ m. We roll off the acceleration below f_{crit} , so that $S_{\text{na}}(f \leq f_{\text{crit}}) = S_{\text{na}}(f_{\text{crit}})$. In reality, the LISA noise will probably not follow this simple power law all the way down to our choice of f_{crit} , but will begin to rise at a higher frequency.

3. RESULTS

3.1. Physical Properties & Numbers

3.1.1. Numbers: Model A

In Table 1 we present the number of double compact objects predicted for the present time in our Galaxy. The numbers are given for each Galactic component (i.e., disk, bulge, halo) and the contribution of each type of double compact object binary (i.e., NS-NS, BH-NS, BH-BH) is shown. Additionally to the total number of Galactic double compact objects, we list the number of LISA systems. The LISA systems are the binaries that are most likely to have right properties (see eq. 16) to produce a GR signal in the frequency sensitivity band of LISA. In the following we discuss only results for Model A.

As a whole, the Galactic population of double compact object is dominated by disk (66%) and bulge (30%) systems with a small contribution of halo binaries (4%). This reflects the fact that the number of double compact objects is proportional (for a given evolutionary model) to the stellar mass. Since the disk stellar mass is the largest, the disk population of double compact objects is the most numerous. The bulge stellar mass is about half that of the disk, and the number of bulge double compact object binaries is about half of those in the disk. Similarly for the halo. The numbers are not exactly proportional to the initial stellar mass, since the star formation history and initial conditions (e.g., metallicity) alter the formation efficiency. However, these factors are only secondary in comparison to the dependence of double compact object number on the initial stellar mass.

Within the entire double compact object population, BH-BH systems are dominant (67%), with a significant contribution of NS-NS binaries (28%) and the very small relative number of BH-NS systems (5%). The dominance of BH-BH systems is a consequence of the natal kick model employed, i.e., neutron stars are given full kicks, while black holes (due to formation through either partial fall back or direct stellar collapse) receive smaller or no natal kicks at all. This naturally leads to the survival of BH-BH progenitors (especially those on wide orbits) while the vast majority of NS-NS progenitors are disrupted at either the first or second supernova. Progenitors of BH-NS systems are depleted by two processes. First, the large difference in initial mass of two stars (one very massive to make a BH, and the other of lower mass to produce a NS) tends to lead to a merger in the first RLOF episode early in the evolutionary history of such systems. Second, many systems that survive the first RLOF and subsequent formation of a BH, are disrupted by the natal kick during NS formation.

The population of potential LISA sources is quite different than the entire double compact object population. These are the binaries that were formed at close orbits ($P_{\text{orb}} \lesssim 150$ day), and they constitute only a small fraction of the entire population. The NS-NS LISA binaries are 30% of entire Galactic NS-NS population, while it is 7% for BH-NS and only 1% for BH-BH LISA binaries. The decreasing contribution to the LISA population is due to the fact that many more BH-BH systems were allowed to form on rather wide orbits while wide NS-NS binaries were mostly disrupted by natal kicks, as explained above.

The bias towards disk binaries is even more pronounced for the LISA binaries, in that they mostly originate from the disk (78.8%), with significant contribution from the bulge (20.9%) and a negligible number of the systems in halo (0.3%). This increased disk contribution (relative to the entire double compact object population) is an effect of star formation history. Basically, the LISA group consists of such close binaries that they can merge (and disappear from the population) in less than the age of Galaxy. Since the disk is forming stars in an approximately constant manner there is a constant supply of close binaries, while for bulge and halo that have formed all the stars ~ 10 Gyr ago, many close binaries have already merged.

The most striking feature of LISA population is that it mostly consists of NS-NS binaries (90.4%), with only minor contributions from BH-NS (3.7%) and BH-BH (5.9%) systems. For single star evolution with the adopted IMF it is predicted that NSs outnumber BHs by about factor of 5.

Binary evolution further affects the numbers. Double neutron star progenitors are more likely to avoid mergers in RLOF episodes as the two stars in the binary are close in mass, while for BH-BH progenitors as mentioned above it is likely that dynamical instability develops and a progenitor system enters a common envelope that may lead to a merger. Since, in the case of the LISA population, we consider only the population of progenitors that start in close orbits (to form the very close double compact objects) kicks that are higher for neutron stars

⁰The IMF slope is -2.7, neutron star progenitor mass range in our single star model is $7.5 - 21 M_{\odot}$ while for black holes it is $21 - 150 M_{\odot}$ (see Belczynski et al. 2008; their Fig. 1).

are not an important factor that could allow binaries with BHs to outnumber NS-NS systems. Additionally, neutron stars with less massive (and smaller in size) progenitors can start their evolution from closer orbits and therefore form double compact object binaries on closer orbits as compared with BH progenitors. The dominance of NS-NS binaries over BH-NS and BH-BH systems was already noted in our models for a different Galactic population (binaries with even closer orbits) of double compact object mergers (Belczynski et al. 2007).

3.1.2. Physical Properties: Model A

In Figure 1 we show the characteristic properties of double compact objects that may be detected by LISA. The panels show distributions of orbital frequencies ($f_{\text{orb}} = 1/P_{\text{orb}}$; top), eccentricities (middle) and chirp masses (bottom) of LISA double compact objects. There is a subtle difference between the distributions presented here and in various papers for double compact objects (e.g., Belczynski et al. 2007). Mostly the results for double compact objects are calculated for a uniform population of stars and are presented at the double compact object formation or for the population that will merge within some specific time (e.g., Hubble time). The distributions presented here, however similar they may appear, are none of the above. First, our distributions originate from the mixture of stellar populations used to represent the Galaxy (different metallicities and star formation histories). Second, double compact objects form at a given time (set primarily by a specific star formation history) and then are evolved to the present time. During the evolution each compact object is subject to the emission of GR and its orbit (period and eccentricity) is decaying. Finally, we only select and present double compact objects that satisfy Eq. 16.

The frequency distributions are similar for all types of double compact objects and they span a wide range: $f_{\text{orb}} \sim 10^{-3} - 10^{-7}$ Hz ($P_{\text{orb}} \sim 0.02 - 150$ day) and peak at $f_{\text{orb}} \sim 5 \times 10^{-5}$ Hz ($P_{\text{orb}} \sim 0.5$ day). The shape of the frequency distributions may be understood in the following way. For frequencies lower than $f_{\text{crit}} = 5 \times 10^{-5}$ Hz (or large orbital periods) the number of systems is falling off since the eccentricity of a given system needs to increase with decreasing frequency in order for the system to be detectable (see eq. 16). However, the number of systems with higher and higher eccentricities is gradually decreasing despite the fact that eccentricity distributions are rather flat (see below). For high frequencies ($f_{\text{orb}} > f_{\text{crit}}$) the systems are so tight ($P_{\text{orb}} < 0.5$ day for a circular binary) that the orbital decay is very fast due to a very effective emission of GR. The tighter the system (higher f_{orb}), the shorter time it takes for a system to merge causing a depletion in number with an increasing GR frequency.

The eccentricity distributions are rather flat for all double compact objects. For NS-NS and BH-NS binaries the distributions are slightly skewed toward the high e -values, while it is just the opposite for BH-BH systems. The eccentricity distribution is the direct result of the second supernova asymmetry; both natal kick and mass loss from the exploding star. Since systems with neutron stars receive, on average, larger second kick the NS-NS and BH-NS binaries are more eccentric than BH-BH systems. The rather high fraction ($\sim 40\%$) of BH-BH systems with small eccentricities ($e < 0.2$) is either the result of direct collapse

of a star to a BH or significant fall back of material during the second supernova explosion.

Chirp mass ($M_{\text{chirp}} = (M_1 M_2)^{0.6} / (M_1 + M_2)^{0.2}$) distributions are very different for the three subclasses of double compact objects. The NS-NS binaries have a chirp mass distribution that peaks at $M_{\text{chirp}} \sim 1.2 M_{\odot}$ with a tail that extends to $M_{\text{chirp}} \sim 2 M_{\odot}$. Systems with black holes have much flatter distributions but spanning a wide range of chirp masses; $M_{\text{chirp}} \sim 1.5 - 4 M_{\odot}$ for BH-NS and $M_{\text{chirp}} \sim 2.5 - 9 M_{\odot}$ for BH-BH binaries. Most neutrons stars form with mass $1.2 - 1.4 M_{\odot}$ either through electron capture supernova, or through regular FeNi core collapse (stars with initial mass $M_{\text{zams}} \lesssim 18 M_{\odot}$; Timmes et al. 1996) and they explain the peak at low chirp mass. The accretion onto the first born neutron star was calculated (Belczynski et al. 2007) to be rather small ($\sim 0.05 - 0.1 M_{\odot}$) and just high enough to mildly recycle the pulsar. The mild recycling and the range of neutron star masses predicted here agrees rather well with the observed masses in the known NS-NS systems, all of which are found with $1 - 1.5 M_{\odot}$ (e.g., Lorimer 2005). Only very few neutron stars are predicted with higher masses in the NS-NS population since they are formed from initially massive stars and in a narrow range ($18 \lesssim M_{\text{zams}} \lesssim 21 M_{\odot}$) and so they are not predicted to show up in such a small observed sample (8 systems). For BH-NS and BH-BH systems the distributions are mostly shaped by the mass distribution of black holes. The black holes are predicted within the very wide mass range and that makes the chirp mass distributions wide as well, wider for BH-BH systems and somewhat narrower for BH-NS binaries since for these systems one component is always of a low mass ($\sim 1.3 M_{\odot}$). The lowest mass black holes are found just over $2.5 M_{\odot}$ as we adopted such a value as the maximum neutron star mass and they can be as massive as $\sim 15 M_{\odot}$ for disk and bulge (high metallicity) while they can reach even higher masses $\sim 30 M_{\odot}$ for halo population (low metallicity environment). The very high masses of black holes predicted for the halo are the result of the evolution of stars in a low metallicity environment; stars lose much less mass in winds so there is a much larger mass reservoir for compact object formation as compared to high metallicity models (disk and bulge). Here our predictions are also consistent with the observations of binaries hosting confirmed stellar mass black holes. Stellar mass black holes are found in our Galaxy with masses reaching $\sim 15 M_{\odot}$ (e.g., Casares 2006), while in the external galaxy IC-10 with a low metallicity environment a black hole was found with mass $\sim 24 - 33 M_{\odot}$ (Prestwich et al. 2007; Silverman & Filipenko 2008). Note that although we predict such high mass black holes in the halo of our Galaxy there are very few of them, and they do not have very massive companions as the highest chirp masses for BH-BH binaries are $M_{\text{chirp}} < 10 M_{\odot}$. The system in IC-10 was predicted to form a BH-BH binary with chirp mass of $M_{\text{chirp}} \sim 20 M_{\odot}$ but with a rather short coalescence time $\sim 2 - 3$ Gyr (Bulik, Belczynski & Prestwich 2008). Therefore, if such a system has formed in the halo of our Galaxy it has most probably merged by now.

3.1.3. Model B

In Table 1 we also list the numbers of double compact object binaries for our Model B, in which we do not allow for survival through common envelope with Hertzsprung gap donors.

For entire population the most affected systems are BH-NS systems (reduction by factor of ~ 2), and BH-BH binaries (reduction by ~ 1.4) while NS-NS binaries are affected the least (reduction by ~ 1.2). All these changes are not very large especially in the light of the limited prospects for detecting the majority of these binaries; since the majority of these systems are too wide to ever make it to the LISA frequency range or to merge within a Hubble time to be detected by ground-based detectors like LIGO or VIRGO.

For LISA binaries, the differences are much more pronounced. BH-BH binaries are reduced by factor ~ 250 , BH-NS by factor ~ 10 , while NS-NS by factor ~ 1.5 . The progenitors of close LISA binaries are subject to one or more common envelope events and thus they are greatly affected in this model. In particular, progenitors of binaries with black holes that start their evolution with stars of rather unequal masses are subject to evolving through the common envelope and merging. Basically, it is predicted that if, in fact, there is no survival in a common envelope with Hertzsprung gap donors, then there are almost no BH-BH and BH-NS binaries in our Galaxy within the LISA frequency band. In such a case, out of all double compact objects, only close NS-NS binaries would have a chance to show up in the LISA datastream.

Although the number of potential LISA NS-NS binaries is reduced, it is still quite significant ($\sim 8 \times 10^4$). The physical properties of LISA NS-NS binaries in Model B are not much different from Model A, so it is expected that the GR signal will have similar shape and will only be reduced in strength. For comparison with model A we show physical properties of model B potential LISA double compact objects in Figure 2.

3.2. Spatial Distribution

We present the cumulative distribution (after propagation) of distances of DCO from the Earth in Figure 3. The distributions corresponding to DCO originating in the three components: disk, bulge and halo are shown separately. The distribution of binaries in each component is determined by their initial positions and velocities, as well as the distribution of merger times. If the mean merger time is longer than a few orbits than the distribution of binaries may relax to a state determined only by the distribution of energies and angular momenta of DCOs in each component. Since the merger time distribution is typically $\propto t^{-1}$ the mean merger time is typically half the age of the Galaxy, i.e. much longer than a few orbits in the Galactic potential. Thus the distribution of DCOs in each component is close to equilibrium in the potential. The energies and angular momenta of the binaries are determined by their initial positions and velocities. The halo originating binaries are born far from the Galactic center and their potential energies are large, the kick velocities may increase them above zero and unbind some of them. There is no preferred direction of the angular momentum for these binaries. Thus their distribution is nearly isotropic around the Galactic center. The density profile is almost flat in the central 10 – 20kpc, and then it starts to fall off. From

the Earth we see their number increasing initially as the distance cube and then the dependence flattens out beyond ≈ 20 kpc. The bulge originating binaries have low potential energies, and adding the kick velocities unbinds only a small fraction of them. Their angular momenta are not large since they are all born close to the Galactic center. Thus they remain concentrated in the bulge. The cumulative number as seen from the Earth is determined by the small number of escaping binaries at distances smaller than ≈ 7 kpc, than it goes up very fast when the Galactic center is probed. Above ≈ 9 kpc the distribution flattens out. The binaries originating in the disk present a more complicated case. Their potential energies have a wide distribution, depending on the distance from the Galactic center, however, the disk itself is a potential well. The addition of kick velocities increases the potential energy. The angular momenta of the disk binaries on their Galactic orbits are initially aligned perpendicularly to the disk plane, yet after the propagation the kick velocities tend to partially isotropize them. Thus, they form a disk with scale height of ≈ 0.2 kpc and the radius of ≈ 15 kpc. The distribution as seen from the Earth initially probes this disk for small distances and the number of DCOs increases with a cube of distance, yet this dependence is only seen for the closest few tens of binaries. For the distances from a few hundred pc to ≈ 10 kpc the number of DCOs increases roughly as the distance squared because a flat distribution of sources is probed. For larger distances ($\gtrsim 15$ kpc) the dependence flattens out because we reach the end of the disk. The disk originating binaries are the main contribution to the sources observable from Earth for all distances.

3.3. Gravitational Radiation Signature

The gravitational wave signature for a 1-year observation has been calculated for both models. The population has been separated into the different binary types in Figure 4 and compared with a realization of the LISA noise. In Figure 5, we separate out the different components of the Galaxy. From this, we see that in both models, the disk binaries are the only ones to rise above the LISA noise.

We are interested in determining if LISA observations can distinguish between these two models. Although we do not have an algorithm for identifying eccentric stellar mass compact object binaries in the LISA data stream, we can still approximate the outcome of such an algorithm. It is possible that the coherent signals from the many harmonics of eccentric binaries can be summed to boost a signal above the LISA noise even though each individual harmonic lies buried in the noise (Benacquista 2001, 2002, Larson & Hellings 2009). We assume that some sort of matched-filter will be used, and so we can describe the detection statistic as the signal-to-noise ratio (ρ) defined as (Wainstein & Zubakov, 1962):

$$\rho^2 = 4 \int_0^\infty \frac{|\tilde{h}(f)h(f)|}{S_n(f)} df, \quad (20)$$

where $S_n(f)$ is the one-sided noise power spectral density. For the purposes of calculating the detection statistic, we use the discrete analog of Equation 20 given by Schutz

(1997):

$$\rho^2 = \sum_{k=0}^{N-1} \frac{|\tilde{h}_k|^2}{S_k} \quad (21)$$

and take S_k to be the running average over 1000 bins of the power spectrum of the LISA noise given by Equation 19 combined with a total Galactic white dwarf binary population generated using StarTrack (Ruiter et al. 2009a,b) and the DCO population considered in this work. We calculate $\tilde{h}_k$ for each binary in this work using the LISA signal simulator described in Section 2.3.

If the foreground noise from the confusion-limited signal from the Galactic population of close white dwarf binaries were Gaussian, we could simply assign a reasonable threshold value of ρ as a criterion for detection. However, the foreground is not Gaussian (Timpano et al. 2006). Racine & Cutler (2007) have studied the impact that the non-Gaussianity of the WD-WD foreground has on the detection of extreme mass ratio inspirals (EMRIs) and super-massive black hole inspirals (SMBHs). Their conclusion is that the threshold criterion is essentially unchanged after the loudest white dwarf binary signals have been removed from the foreground signals. Their result can be explained by the fact that the time-scale of the evolution of EMRIs, SMBHs, and WD-WDs are all different. Although the evolutionary time-scales of stellar mass compact object binaries are not significantly different from those for white dwarf binaries, white dwarf binaries are almost exclusively circularized in the LISA band and nearly all compact object binaries analyzed in this work are eccentric. Thus, their waveforms differ significantly in the time domain. This can be used to distinguish between the two populations, and we assume that the results of Racine and Cutler (2007) can be applied here.

Finally, we note that the imposition of a threshold signal-to-noise ratio calculated according to Equation 21 can result in detection of binaries for which no individual harmonic has a signal-to-noise ratio above 1. Matched filtering algorithms have already been demonstrated that can successfully detect SMBH signals within mock LISA data (Babak et al. 2008) even though the signal never exceeds the instrument and white dwarf binary confusion noise at any frequency (see Fig 1 of Arnaud et al. 2007).

Consequently, we impose a threshold of $\rho \geq 10$ as the criterion for detection. Using this estimate, we find that there are 11 resolvable double compact object binaries in model A and only 2 in model B. From Figure 4, we see that the bulk of the potentially detectable very low frequency sources in model A are BH-BH binaries and that these are dramatically suppressed in model B. This is also borne out by the estimates of resolvable binaries where the number of resolvable BH-BH binaries drops from 5 in model A to zero in model B. Thus, detection of high-mass, low-frequency binaries can be used to distinguish between these two models. In model A there is a relatively large number of BH-BH binaries at low frequencies (at 10^{-4} Hz there is a comparable number of NS-NS and BH-BH systems; see Fig. 1). Since at these frequencies binaries have relatively large orbital separations, neutron stars that are of much lower mass than black holes do not generate a detectable GR signal, while massive black hole systems do. For lower frequencies ($\lesssim 10^{-5}$ Hz) corresponding to yet

larger orbital separations, even the most massive BH-BH binaries do not generate detectable signal, although they dominate over other double compact objects. The properties of the potentially resolvable binaries are presented in Table 2 and shown in Figures 1 and 2.

4. COMPARISON WITH PREVIOUS WORK

The number of detectable compact object binaries that we found in this analysis is somewhat smaller than the 42 detectable systems predicted by Nelemans et al. (2001). However, this can be reasonably explained by the differences in the overall population numbers between the two simulations. The total number of compact object binaries predicted in our simulation ranges between 1.0×10^6 for model B and 1.4×10^6 for model A, while Nelemans et al. predict 4.0×10^6 such systems. Thus, our prediction of ~ 11 systems is comparable with Nelemans et al. It should be noted, however, that our estimate is conservative as we have assumed no prior removal of any WD-WD signals from the calculated value of S_k , and that we have assumed an observation time of 1 year. In reality, we expect some removal of the loud WD-WD signals and the nominal mission lifetime of LISA is 3 years. On the other hand, we have not included any penalties incurred by the larger parameter space needed for the analysis of eccentric systems.

5. CONCLUSIONS

We have calculated, in a self-consistent way, the gravitational radiation signal of Galactic double compact object binaries. The realistic model for Galactic stars was employed, then massive stars were followed in their evolution to form neutron stars and black holes, some of which were formed in binaries. The current position of the binaries with neutron stars and black holes relative to Earth was obtained and integrated gravitational radiation signal was predicted for the LISA detector.

It was found that several double compact object binaries can be detected and resolved in a 1-year observation with an instrument like LISA. In model A, we find comparable numbers of BH-BH and NS-NS binaries, while no BH-BH binaries are observed in model B. This prediction, if true, would put a new light on double compact object studies as at present we have only detected 9 NS-NS systems. Double compact objects with a black hole are yet to be discovered. LISA could potentially detect several of these binaries. Starting with NS-NS systems, these potential detections could answer a number of questions. Do nearly all NS-NS systems host low mass neutrons stars ($\sim 1.35 M_\odot$) as it seems to appear in a known sample discovered through radio-pulsar surveys or is it only some observational bias? Galactic merger rates seem to appear consistent with the notion that NS-NS are short GRB progenitors. But would this be confirmed by a broader sample detected by LISA? Detection of any double compact objects with black holes would be a feat in and of itself. Although a number of these systems are expected from evolutionary calculations, these are usually burdened with rather large uncertainties. Detections of BH-BH binaries would provide very valuable tests for evolutionary codes that are used in number of studies. Knowledge of black hole numbers and their basic properties can put direct constraints on evolution of the

most massive stars, their progenitors. In particular, one such example was presented in our study. Depending on a common envelope model the number of BH-BH systems that could be detected with LISA vanishes. So just by their presence (or lack of thereof) in the signal one could test an evolutionary phase that is virtually undetectable due to its shortness ($\sim 10^3 - 10^4$ yr). And it is interesting to note, that although the common envelope was proposed more than 30 years ago (Paczynski 1976), there is still a surprising lack of understanding of the outcome—and the outcome is crucial for formation of Type Ia progenitors (close white dwarf binaries) or any type of X-ray binaries that are currently observed even in external galaxies by *Chandra* or *XMM*, just to name a few examples.

Although, double compact objects will be most probably discovered first in gravitational radiation with ground-based interferometers (e.g., LIGO, VIRGO) they will be detected only in the inspiral and merger phases, preventing a measurement of their orbital parameters (e.g., separations and eccentricities). Additionally, the merger rates are too small to expect detections in the Galaxy, and the

observations will be made for a wide spectrum of local environments in which mergers are taking place, making it harder to connect the result to a particular evolutionary model that by nature depends on local conditions (e.g., metallicity, IMF or age of population). In this context, LISA may provide a potentially very unique sample of double compact object binaries in the most studied environment of our own Galaxy and in the same providing the most robust constraints on evolution of massive stars.

ACKNOWLEDGMENTS

We would like to thank two anonymous referees for a number of helpful comments, Joe Romano for providing a LISA noise generation code and Ashley Ruiter for providing WD-WD foreground noise sources. KB and TB acknowledge the support from the Polish Ministry of Science and Higher Education (MSHE) grant N N203 302835 (2008-2011). MJB acknowledges the support of NASA Grant NNX08AB74G and the Center for Gravitational Wave Astronomy. MJB and KB both acknowledge the support and hospitality of the Aspen Center for Physics.

REFERENCES

- Abt, H. A. 1983, *ARA&A*, 21, 343
- Arnaud, K., et al. 2007, *Classical and Quantum Gravity*, 24, S551
- Babak, S., et al. 2008, *Classical and Quantum Gravity*, 25, 114037
- Benacquista, M., DeGoes, J., & Lunder, D. 2004, *Classical and Quantum Gravity*, 21, 509
- Benacquista, M. 2002, *Classical and Quantum Gravity*, 19, 1297
- Benacquista, M. 2001, 20th Texas Symposium on Relativistic Astrophysics, 586, 793
- Belczynski, K., & Bulik 1999, *A&A*, 346, 91
- Belczynski, K., Kalogera, V., & Bulik, T. 2002a, *ApJ*, 572, 407
- Belczynski, K., Taam, R., Kalogera, V., Rasio, F., & Bulik, T. 2007, *ApJ*, 662, 504
- Belczynski, K., Kalogera, V., Rasio, F., Taam, R., Zezas, A., Bulik, T., Maccarone, T., & Ivanova, N. 2008a, *ApJS*, 174, 223
- Bulik, T., Belczynski, K., & Prestwich, A. 2008, *ApJ*, submitted (arXiv:0803.3516)
- Bethe, H.A., & Brown, G.E. 1998, *ApJ*, 506, 780
- Bloom, J., Sigurdsson, S., & Pols, O. 1999, *MNRAS*, 305, 763
- Casares, J. 2006, IAU Symposium 238: "Black Holes: From Stars to Galaxies -Across the Range of Masses", in press (astro-ph/0612312)
- Cornish, N.J. 2001, *Phys. Rev. D*, 65, 022004
- Cutler, C. 1998, *Phys. Rev. D*, 57, 7089
- Dewi, J.D.M., & Pols, O.R. 2003, *MNRAS*, 344, 629
- Duquennoy, A., & Mayor, M. 1991, *A&A*, 248, 485
- Farmer, A.J., & Phinney, E.S. 2003, *MNRAS*, 346, 1197
- Fryer, C., Woosley, S., & Hartmann, D. 1999, *ApJ*, 526, 152
- Fryer, C., & Kalogera, V. 2001, *ApJ*, 554, 548
- Grindlay, J., Portegies Zwart, S., McMillan, S. 2006, *Nature*, 2, 116
- Gultekin, K., Miller, M., & Hamilton, D. 2004, *ApJ*, 616, 221
- Hils, D., Bender, P.L., & Webbink R.F. 1990, *ApJ*, 360, 75
- Heggie, D. C. 1975, *MNRAS*, 173, 729
- Hobbs, G., Lorimer, D.R., Lyne, A.G., & Kramer, M. 2005, *MNRAS*, 360, 974
- Hughes, S. 2006, *Laser Interferometer Space Antenna: 6th International LISA Symposium*, 873, 13
- Hurley, J. R., Pols, O. R., & Tout, C. A. 2000, *MNRAS*, 315, 543
- Hurley, J. R., Tout, C. A., & Pols, O. R. 2002, *MNRAS*, 329, 897
- Ivanova, N., Heinke, C., Rasio, F., Belczynski, K., & Fregeau, J. 2008, *MNRAS*, 386, 553
- Kasian, L. 2008, 40 YEARS OF PULSARS: Millisecond Pulsars, Magnetars and More. AIP Conference Proceedings, 983, 485
- Kim, C., Kalogera, V., & Lorimer, D. 2006, *New Ast. Rev* (astro-ph/0608280)
- Kroupa, P., Tout, C.A., & Gilmore, G. 1993, *MNRAS*, 262, 545
- Kroupa, P., & Weidner, C. 2003, *ApJ*, 598, 1076
- Kulkarni, S., Hut, P., & McMillan, S. 1993, *Nature*, 364, 421
- Larson, S., & Hellings, R. 2009, *BAAS*, 41, 340.
- Levine, A., Rappaport, S.A., & Zojcheski, G. 2000, 541, L194
- Lipunov, V., Postnov, K., & Prokhorov, M. 1997, *MNRAS*, 288, 245
- Lorimer, D. 2005, *Living Rev. Relativity*, 8, 7 (www.livingreviews.org/lrr-2005-7)
- Nelemans, G., Yungelson, L., & Portegies Zwart, S. 2001, *A&A*, 375, 890
- Nelemans, G., Yungelson, L., & Portegies Zwart S. 2004, *MNRAS*, 349, 181
- O'Leary, R. M., Rasio, F. A., Fregeau, J. M., Ivanova, N., O'Shaughnessy, R. 2006, *ApJ*, 637, 937
- Paczynski, B. 1976, in *Structure and Evolution of Close Binary Systems; Proceedings of the Symposium*, Cambridge, England, Edited by P. Eggleton, S. Mitton, and J. Whelan. Dordrecht, D. Reidel Publishing Co. (IAU Symposium No. 73), 1976., p.75
- Peters, P., & Mathews, J. 1963, *Phys. Rev.*, 131, 434
- Peters, P. 1964, *Phys. Rev.* 136, 1224
- Pfahl, E., Podsiadlowski, P., & Rappaport, S. 2005, *ApJ*, 628, 343
- Phinney, S. 1991, *ApJ*, 380, L17
- Pierro, V., Pinto, I., Spallicci, A., Laserra, E., & Recano, F. 2001, *MNRAS*, 325, 358
- Podsiadlowski, P., Langer, N., Poelarends, A.J.T., Rappaport, S., Heger, A., & Pfahl, E.D. 2004, *ApJ*, 612, 1044
- Portegies Zwart, S., & Yungelson, L. 1998, *A&A*, 332, 173
- Portegies Zwart, S., & McMillan, S. 2000, in *Black Holes in Binaries and Galactic Nuclei. Proceedings of the ESO Workshop*. Lex Kaper, Edward P. J. van den Heuvel, Patrick A. Woudt (eds.), p. 252. Springer.
- Prestwich, A., et al. 2007, *ApJ*, 669, L21
- Racine, E. & Cutler, C. 2007, *Phys. Rev. D*, 76, 124033
- Rubbo, L., Cornish, N., & Poujade, O. 2004 *Phys. Rev. D*, 69, 082003
- Ruiter, A., Belczynski, K., Benacquista, M., & Larson, S. 2009a, *ApJ*, submitted (arXiv:0705.3272)
- Ruiter, A., Belczynski, K., Benacquista, M., & Holley-Bockelmann, K. 2009b, *ApJ*, 693, 383
- Sadowski, A., Belczynski, K., Bulik, T., Ivanova, N., Rasio, F., & O'Shaughnessy, R. 2008, *ApJ*, 676, 1162
- Schutz, B. 1997, in *Fundamental Physics in Space, Proceedings of the Alpbach Summer School*, Alpbach, Tyrol, Austria, 22-31 July, 1997, A. Wilson (ed), European Space Agency Publication, p. 265
- Silverman, J., Filipenko, A., 2008, arXiv:0802.2716
- Taam, R. E., & Sandquist, E. L. 2000, *ARA&A*, 38, 113
- Timmes, F.X., Woosley, S.E., & Weaver, T.A. 1996, *ApJ*, 457, 834
- Timpano, S.E., Rubbo, L.J., & Cornish, N.J. 2006, *Phys. Rev. D*, 73, 122001
- Tutukov, A., & Yungelson, L. 1994, *MNRAS*, 268, 871
- Vecchio, A. & Wickham, E.D.L., 2004, *Phys. Rev. D*, 70, 082002
- Voss, R., & Tauris, T. 2003, *MNRAS*, 342, 1169
- Wainstein, L.A. & Zubakov, V.D. 1962, "Extraction of Signals from Noise", Prentice-Hall
- Webbink, R. 1984, *ApJ*, 277, 355
- Wellstein, S., Langer, N., & Braun, H. 2001, *A&A*, 369, 939

TABLE 1
NUMBER OF DOUBLE COMPACT BINARIES IN THE GALAXY^a

	NS-NS	BH-NS	BH-BH
ALL: disk	281091 – 238245	45075 – 17729	621812 – 424501
ALL: bulge	58658 – 52520	10347 – 4388	147260 – 106170
ALL: halo	6351 – 6342	2845 – 2776	37681 – 36373
ALL: total	346100 – 297107	58267 – 24893	806753 – 567044
LISA: disk	92253 – 58633	4029 – 371	6903 – 6
LISA: bulge	12853 – 8700	382 – 48	417 – 0
LISA: halo	298 – 291	50 – 39	36 – 25
LISA: total	105404 – 67624	4461 – 458	7356 – 31

^aNumbers are given for two models: Model A – Model B. ALL – all binaries in a given category. LISA – binaries with orbital periods and eccentricities that are likely to contribute to GR signal in LISA frequency band.

TABLE 2
PROPERTIES OF RESOLVABLE DOUBLE COMPACT BINARIES IN THE GALAXY

Model	Type	M_{chirp} ($M_{\odot}$)	f_{orb} (mHz)	e	ρ
A	BH-BH	5.35	0.284	0.038	22.7
A	BH-BH	5.77	0.455	0.005	22.1
A	BH-BH	4.01	0.448	0.011	17.3
A	BH-BH	7.69	0.097	0.006	15.2
A	BH-BH	7.46	0.279	0.006	11.5
A	NS-NS	1.16	1.449	0.167	53.4
A	NS-NS	1.24	0.187	0.884	32.1
A	NS-NS	1.16	0.760	0.346	17.3
A	NS-NS	1.15	0.956	0.044	13.2
A	NS-NS	1.16	0.843	0.106	12.2
A	NS-NS	1.15	0.990	0.069	10.0
B	NS-NS	1.16	0.760	0.346	17.3
B	NS-NS	1.15	0.990	0.069	10.0




Fig. 1.— Potential LISA Galactic population of double compact objects (Model A). The top panel shows the distribution of orbital frequency, middle panel the distribution of eccentricities, while the bottom panel has the distribution of chirp masses. Note the dominance of NS-NS binaries. At the top of each panel we show Galactic resolved double compact objects. Since this population is very small (6 NS-NS and 5 BH-BH binaries) we show all individual resolved sources (details in Table 2).




Fig. 2.— Same as Fig. 1 but for model B. The NS-NS dominance is even more pronounced in this model. Note that there are only 2 resolved sources and they are both NS-NS binaries.




Fig. 3.— The cumulative distribution of the current distances of simulated binaries from the Earth (model A). The distances are obtained after propagation in the Galactic potential; the binaries have spread out from their birth sites due to natal kicks. We show the three Galactic components separately. Note that the Galactic center is at a distance of ~ 8 kpc from Earth.




Fig. 4.— Strain spectral densities of the double compact object binaries separated by binary type and compared with the full spectrum containing the double compact objects, the WD-WD signal and the LISA noise. The line is a running average over the WD-WD and LISA noise spectrum. The left panel shows model A and the right panel shows model B. Note the strong suppression of the BH-BH binaries in model B, and the absence of any detectable signals below ~ 0.4 mHz.




Fig. 5.— Strain spectral densities of the double compact object binaries separated by Galaxy component and compared with the full spectrum containing the double compact objects, the WD-WD signal and the LISA noise. The line is a running average over the WD-WD and LISA noise spectrum. The left panel shows model A and the right panel shows model B. Note the dominance of the disk binaries in both models, but the significant absence of low frequency binaries in model B.